

Appendix D

Discrepancy Score Tables

Table D.1. Cumulative Percentages of the Normative Samples Obtaining PHC>MHC T-Score Discrepancies

Amount of Discrepancy	Age-Based Sample				Age-Stratified Sample		
	18-24	25-44	45-64	≥65	Overall	Female	Male
≥28	0.5	0.0	1.0	0.5	0.4	0.8	0.0
27	0.5	0.0	1.0	0.5	0.6	1.2	0.0
26	0.5	0.5	1.0	0.5	1.2	2.4	0.0
25	0.5	0.5	1.0	0.5	1.4	2.7	0.8
24	1.0	1.5	1.0	0.5	1.4	2.7	0.8
23	1.0	1.5	1.5	0.5	1.8	2.7	0.8
22	1.0	3.0	2.0	1.0	2.4	3.9	0.8
21	1.0	3.0	2.0	1.0	2.4	4.3	0.8
20	1.0	3.0	2.0	1.5	2.6	4.3	1.2
19	1.0	4.0	3.0	1.5	3.0	4.7	1.6
18	1.0	4.5	3.5	1.5	3.6	5.5	2.4
17	2.5	4.5	4.5	1.5	4.0	5.5	3.7
16	4.5	5.0	4.5	3.0	4.6	5.5	4.1
15	6.0	5.5	5.0	4.5	5.0	5.5	4.5
14	6.5	6.0	5.5	6.0	5.2	6.7	4.9
13	7.0	6.5	6.5	7.0	6.8	8.2	5.7
12	8.5	7.0	8.0	8.0	8.2	10.2	6.9
11	8.5	9.5	8.5	8.0	10.2	13.7	8.6
10	12.0	12.0	10.0	8.5	12.0	14.9	11.0
9	13.5	13.0	11.5	10.5	14.4	17.6	13.5
8	17.5	15.5	14.5	12.5	16.6	19.2	16.7
7	20.0	17.5	16.5	15.5	19.2	22.7	20.8
6	24.5	21.0	20.0	20.0	23.0	25.1	24.9
5	26.0	24.0	22.0	24.5	26.4	29.0	26.9
4	31.0	29.0	27.0	27.5	29.4	32.5	30.6
3	37.5	35.5	31.5	35.0	32.4	36.1	35.9
2	41.5	42.5	35.5	44.0	36.8	38.8	41.2
1	44.0	47.5	42.5	47.0	42.2	44.3	46.9
0 ^a	7.0	7.5	6.0	6.0	5.8	5.5	6.2
Mean	-0.2	0.2	0.0	0.1	-0.4	0.0	0.2
SD	8.9	8.5	8.1	7.6	8.9	9.9	8.1
Median	0.0	0.0	-1.0	0.0	-1.0	-1.0	0.0

Note. PHC-MHC discrepancies >10 were obtained by 15% or less of individuals within each normative group. PHC-MHC discrepancies >6 are statistically significant at $p < .10$. This table does not include negative PHC-MHC discrepancies. Mean difference scores, standard deviations, and medians are based on the entire distribution.

^aThis row provides the percentage of individuals in each normative group for whom PHC-MHC composite score discrepancies were equal to zero or <1.

Table D.2. Cumulative Percentages of the Normative Samples Obtaining MHC>PHC T-Score Discrepancies

Amount of Discrepancy	Age-Based Sample				Age-Stratified Sample		
	18-24	25-44	45-64	≥65	Overall	Female	Male
26	0.5	0.0	0.0	0.0	0.2	0.0	0.4
25	1.0	0.0	0.0	0.0	0.6	0.4	0.4
24	1.5	0.0	0.0	0.0	0.6	0.8	0.8
23	2.0	0.0	0.5	0.0	0.8	0.8	0.8
22	2.0	0.0	0.5	0.0	1.2	0.8	0.8
21	2.0	1.0	0.5	1.0	1.2	1.2	1.6
20	2.0	1.0	1.0	1.0	1.6	2.0	1.6
19	2.5	1.5	1.5	1.0	1.8	2.0	1.6
18	3.0	2.5	2.0	1.5	2.2	3.1	2.0
17	3.5	3.0	2.0	2.0	3.0	4.3	2.4
16	6.5	3.5	3.5	2.0	3.6	5.1	2.4
15	6.5	4.5	4.0	2.0	4.4	6.3	2.9
14	7.5	5.0	4.5	2.5	5.2	7.8	4.9
13	8.5	8.0	5.0	3.0	7.6	8.6	6.1
12	8.5	8.5	6.0	5.0	8.6	10.6	6.9
11	9.0	10.0	7.0	5.5	11.6	13.7	8.6
10	9.5	12.0	8.5	8.0	12.4	14.5	10.6
9	13.5	13.5	11.0	10.0	15.4	17.6	11.4
8	14.5	15.5	13.5	13.5	18.4	20.8	14.3
7	17.5	17.5	15.0	19.5	22.0	25.5	17.6
6	23.0	21.0	19.5	22.5	26.6	27.8	20.4
5	24.5	25.5	23.5	26.0	31.2	30.2	24.5
4	32.5	31.5	31.5	31.5	35.8	35.3	30.6
3	38.5	36.5	35.5	36.0	42.8	41.6	35.1
2	44.5	40.0	46.5	40.0	47.0	45.9	43.3
1	49.0	45.0	51.5	47.0	52.0	50.2	46.9
0*	7.0	7.5	6.0	6.0	5.8	5.5	6.2
Mean	0.2	-0.2	0.0	-0.1	0.4	0.0	-0.2
SD	8.9	8.5	8.1	7.6	8.9	9.9	8.1
Median	0.0	0.0	1.0	0.0	1.0	1.0	0.0

Note: MHC-PHC discrepancies >10 were obtained by 15% or less of individuals within each normative group. MHC-PHC discrepancies >6 are statistically significant at $p < .10$. This table does not include negative MHC-PHC discrepancies. Mean difference scores, standard deviations, and medians are based on the entire distribution.

* This row provides the percentage of individuals in each normative group for whom MHC-PHC composite score discrepancies were equal to zero or <1.